
Diverticulitis Diet During Flare Up

*Diverticulitis
Diet During
Flare Up*

*Downloaded from
template-dev.mobaptist.org
by Marisa & Moyer*

UNDERSTANDING DIVERTICULITIS AND THE IMPORTANCE OF DIET DURING A FLARE UP

If you or a loved one has been diagnosed with diverticulitis, you know that managing this condition requires careful attention to digestive health. Diverticulitis occurs when small, bulging pouches (diverticula) that develop in the lining of the digestive system become inflamed or infected. This inflammation can

cause significant discomfort, including abdominal pain, fever, nausea, and changes in bowel habits. While medical treatment is often necessary during an acute episode, one of the most powerful tools you have to manage symptoms and promote healing is your diet. A well-planned diverticulitis diet during flare up can make a substantial difference in how quickly you recover and how comfortable you feel during the process.

The digestive system needs time to rest and repair when

diverticulitis strikes. During an active flare up, the inflamed diverticula are irritated by the passage of food and stool. This is why dietary adjustments are not just helpful—they are essential. By choosing the right foods and avoiding others, you can reduce the workload on your colon, minimize irritation, and support your body's natural healing mechanisms. Understanding the principles of a diverticulitis diet during flare up helps you navigate this challenging period with greater confidence and better outcomes.

What

Happens During a Diverticul itis Flare Up

A diverticulitis flare up is characterized by inflammation and sometimes infection in one or more diverticula. Common symptoms include sudden and persistent pain, typically on the lower left side of the abdomen, along with fever, chills, bloating, and changes in bowel habits such as constipation or diarrhea. The severity of flare ups can range from mild to severe, and dietary management plays a

role at every level. When the colon is inflamed, any food that requires significant digestive effort or leaves substantial residue can aggravate the condition. This is why the diverticulitis diet during flare up focuses on giving the bowel a break while still providing necessary nutrition.

**THE CORE
PRINCIPLE:
RESTING THE
DIGESTIVE SYSTEM**

The primary goal of a diverticulitis diet during flare up is to allow the colon to rest. This means consuming foods that are easy to digest and leave minimal residue in the bowel. The diet typically progresses through several stages as symptoms improve. Initially, your

healthcare provider may recommend a clear liquid diet to give the digestive tract complete rest. As symptoms begin to subside, you can gradually introduce low-fiber foods and eventually transition back to a higher-fiber diet once the flare up has fully resolved.

Stage One: Clear Liquid Diet

During the most acute phase of a diverticulitis flare up, when pain and inflammation are at their peak, a clear liquid diet is often recommended. This

stage typically lasts for a few days and provides hydration and some electrolytes while allowing the colon to rest completely. Acceptable clear liquids include:

- Water and ice chips
- Clear broths (chicken, beef, or vegetable)
- Clear fruit juices without pulp (apple, cranberry, grape)
- Gelatin desserts
- Popsicles without fruit pieces or dairy
- Herbal teas
- Sports drinks for electrolyte balance

Important: Avoid any liquids that are red or purple in color, as they can be mistaken for blood if you experience rectal bleeding. Also,

steer clear of dairy-based liquids and any beverages containing pulp or solid particles. The clear liquid phase of the diverticulitis diet during flare up is temporary and should only be followed under the guidance of your healthcare provider.

Stage Two: Low-Fiber Foods

As your symptoms begin to improve—usually after a few days on clear liquids—you can start introducing low-fiber, soft foods. This transition should be gradual, and you should continue to listen to your body. If

certain foods cause discomfort, eliminate them and return to easier options. Suitable low-fiber foods during this stage of the diverticulitis diet during flare up include:

- White bread, white rice, and plain pasta
- Cooked vegetables without skins or seeds (canned carrots, green beans, potatoes without skin)
- Fruit without skins or seeds (bananas, applesauce, canned peaches)
- Lean proteins such as eggs, fish, and skinless poultry
- Low-fiber cereals like cream of wheat or rice cereals
- Plain yogurt and

cottage cheese (if tolerated)

- Clear soups with soft, low-fiber vegetables

At this stage, it is crucial to avoid any foods that contain seeds, nuts, corn, popcorn, or tough skins. These can potentially lodge in the diverticula and exacerbate inflammation. The low-fiber phase of the diverticulitis diet during flare up typically lasts until all symptoms have resolved, which may take several days to a week.

FOODS TO EMPHASIZE DURING A FLARE UP

Knowing what to eat is just as important as knowing what to avoid.

When following a diverticulitis diet during flare up, certain foods can be particularly beneficial for soothing inflammation and supporting recovery.

Hydrating Fluids

Staying well-hydrated is essential during a diverticulitis flare up. Water helps keep stool soft and supports overall digestive function. Aim to drink at least eight to ten glasses of fluid per day, unless otherwise directed by your doctor. Herbal teas, clear broths, and diluted fruit juices can contribute to your fluid intake while providing variety.

Easy-to-Digest Proteins

Protein is important for tissue repair and immune function. During a flare up, choose lean, tender sources of protein that are easy to digest. Poached or scrambled eggs, baked or broiled fish, and skinless chicken breast are excellent options. Avoid fried or heavily seasoned preparations, as these can be harder on the digestive system.

Low-Fiber Carbohydr

rates

Refined carbohydrates provide energy without placing too much strain on the colon. White bread, white rice, plain crackers, and well-cooked pasta are good choices during the recovery phase of the diverticulitis diet during flare up. These foods are low in residue and can help maintain energy levels while the colon heals.

FOODS TO AVOID DURING A FLARE UP

Equally important to knowing what to eat is understanding which foods to avoid. During an active flare up, certain foods can worsen inflammation, increase pain, and delay healing. Here are the primary categories

of foods to steer clear of when following a diverticulitis diet during flare up:

High-Fiber Foods

While a high-fiber diet is recommended for preventing diverticulitis in the long term, it is counterproductive during an acute flare up. High-fiber foods can irritate the inflamed diverticula and exacerbate symptoms. Avoid whole grains, legumes, nuts, seeds, popcorn, and raw vegetables until the flare up has completely resolved.

Dairy Products (in some cases)

Some people with diverticulitis find that dairy products exacerbate symptoms such as bloating and gas. If you are lactose intolerant or sensitive to dairy, it may be wise to avoid milk, cheese, and ice cream during a flare up. Plain yogurt and cottage cheese are often better tolerated due to their lower lactose content and the presence of probiotics.

Fatty and

Fried Foods

High-fat foods can slow digestion and increase pressure in the colon, potentially worsening symptoms. Avoid fried foods, fatty cuts of meat, rich sauces, and creamy dressings during a diverticulitis flare up. Opt instead for baked, broiled, or steamed preparations with minimal added fat.

Spicy and Highly Seasoned Foods

Spices and strong seasonings can irritate the digestive tract and

should be avoided during the acute phase. Stick to mild flavors and simple preparations. As you recover, you can gradually reintroduce gentle seasonings like salt, pepper, and mild herbs.

Gas-Producing Foods

Some foods are known to cause gas and bloating, which can be particularly uncomfortable during a diverticulitis flare up. Avoid carbonated beverages, cruciferous vegetables (broccoli, cauliflower, cabbage), onions, garlic, and beans until your symptoms have subsided.

Alcohol and Caffeine

Both alcohol and caffeine can irritate the digestive tract and may worsen symptoms. It is best to avoid these beverages during a flare up. Herbal teas, water, and clear broths are safer alternatives that support hydration and healing.

SAMPLE DAILY MEAL PLAN FOR A DIVERTICULITIS FLARE UP

To give you a practical sense of what a diverticulitis diet during flare up might look like, here is a sample menu for the low-fiber phase. Adjust

portions and food choices based on your individual tolerance and your healthcare provider's recommendations.

Breakfast

- 1/2 cup cooked cream of wheat cereal made with water
- 1 scrambled egg
- 1 slice white toast with a thin layer of jelly (seedless)
- 1 cup clear herbal tea

Mid-Morning Snack

- 1 cup applesauce (unsweetened)
- 1 cup water or clear broth

Lunch

- 1/2 cup canned, peeled carrots (well-cooked)
- 3 ounces baked skinless chicken breast
- 1/2 cup white rice
- 1 cup clear broth

Afternoon Snack

- 1 banana (ripe and soft)
- 1 cup water or herbal tea

Dinner

- 3 ounces baked white fish (such as cod or tilapia)
- 1/2 cup mashed potatoes (made with water, not milk or butter)

- 1/2 cup canned green beans (well-cooked)
- 1 cup clear broth

Evening Snack (if desired)

- 1 cup plain gelatin dessert
- 1 cup water

This sample meal plan provides balanced nutrition while keeping fiber intake low and supporting digestive rest. Remember, the diverticulitis diet during flare up should be tailored to your specific needs and tolerance levels.

TRANSITIONING BACK TO A REGULAR DIET

Once your symptoms

have completely resolved, it is important to gradually reintroduce fiber into your diet. A high-fiber diet is one of the most effective strategies for preventing future diverticulitis flare ups. However, the transition should be slow to avoid overwhelming your digestive system.

Step-by-Step Fiber Reintroduction

Start by adding small amounts of soluble fiber, such as oatmeal, bananas, and cooked carrots. After a few days, you can introduce more

substantial sources of fiber, including whole grain bread, brown rice, and cooked vegetables. Finally, add raw vegetables, nuts, seeds, and legumes back into your diet one at a time, monitoring your body's response. Throughout this process, it is crucial to drink plenty of water to help fiber move through your digestive system smoothly.

Long-Term Dietary Habits for Preventio

n

Once you have fully recovered, aim to maintain a diet rich in fiber from fruits, vegetables, whole grains, and legumes. Evidence suggests that a high-fiber diet can reduce the risk of developing diverticulitis and may help prevent recurrent flare ups. Additionally, staying physically active, maintaining a healthy weight, and avoiding smoking are all important lifestyle factors that support colon health.

ADDITIONAL TIPS FOR MANAGING A FLARE UP

Beyond the diverticulitis diet during flare up, there are several other strategies that can

help you manage symptoms and
promote recovery: