

Structure And Function Of The Kidney And Its Parts

Structure And Function Of The Kidney And Its Parts

Downloaded from [template-dev.mobaptist.org](https://www.template-dev.mobaptist.org) by Mathews & Slade

AN INTRODUCTION TO THE BODY’S UNSUNG HERO

Tucked away beneath the left rib cage, the spleen is one of the most versatile and hardworking organs in the human body. Although it often goes unnoticed in day-to-day health discussions, its importance cannot be overstated. For centuries, the spleen was shrouded in mystery—ancient Greeks believed it governed melancholy, while later physicians dismissed it as a vestigial organ. Modern medicine, however, has revealed an astonishing truth: the spleen is a master multitasker, seamlessly blending roles in filtration, immunity, and blood storage. Understanding the **structure and function of the spleen** offers a window into how our bodies maintain balance, fight infection, and recycle vital components. This article will guide you through its remarkable anatomy, its microscopic architecture, and the life-sustaining tasks it performs every minute of every day.

LOCATION AND GROSS ANATOMY

Anatomical Position

The spleen resides in the upper left quadrant of the abdomen, nestled between the stomach, the left kidney, and the diaphragm. It is protected by the 9th, 10th, and 11th ribs, which act as a bony shield. This position is no accident — it allows the spleen to interact closely with the circulatory and lymphatic systems while remaining somewhat insulated from minor trauma. The organ is intraperitoneal, meaning it is suspended within the abdominal cavity by folds of peritoneum, including the gastrosplenic and splenorenal ligaments.

Size and Appearance

In a healthy adult, the spleen is roughly the size of a fist, measuring about 10 to 12 centimeters in length and weighing between 150 and 200 grams. It has a purple-red hue, a soft, spongy consistency, and a shape that is often described as bean-like but more irregular. The organ features two distinct surfaces: a convex diaphragmatic surface that curves against the diaphragm, and a concave visceral surface that contacts the stomach, left kidney, and colon. The splenic hilum, a groove along the visceral surface, serves as the entry and exit point for blood vessels, nerves, and lymphatic channels.

Relations to Other Organs

The spleen does not work in isolation. Its close **anatomical relationship** with the stomach, pancreas, left kidney, and splenic flexure of the colon means that changes in one organ can influence the others. For example, an enlarged spleen (splenomegaly) can compress the stomach, leading to early satiety, or irritate the diaphragm, causing referred shoulder pain. Understanding these spatial relationships is crucial for clinicians assessing abdominal complaints.

THE MICROARCHITECTURE OF THE SPLEEN

To truly appreciate the **structure and function of the spleen**, one must look beyond the naked eye. The internal organization of the spleen is a marvel of biological engineering, divided into distinct compartments that each serve specialized roles.

The Capsule and Trabeculae

The spleen is encased in a fibrous capsule composed of dense connective tissue, which gives the organ its shape and provides some protection. From this capsule, trabeculae—strong, fibrous bands—extend inward, forming a scaffolding that supports the splenic pulp. These trabeculae also carry blood vessels and nerves into the deeper regions of the organ. The capsule contains smooth muscle fibers, which allow the spleen to contract in response to stress or blood loss, squeezing stored blood into the circulation.

Red Pulp

The red pulp is the largest compartment of the spleen, accounting for about 75-80% of its volume. It gets its name from the reddish color imparted by the vast number of red blood cells. Structurally, the red pulp consists of a network of sinusoids (wide, leaky blood vessels) and cords of Billroth—cords of reticular connective tissue filled with macrophages, plasma cells, and blood cells. This region is the primary site for filtration, where old or damaged red blood cells are removed, and iron is recycled. The slow, tortuous flow of blood through the red pulp ensures that every cell is inspected.

White Pulp

Scattered throughout the spleen like islands in a sea of red pulp, the white pulp is the organ’s immune command center. It is composed of lymphoid tissue, primarily B-cells and T-cells, arranged around central arteries. These aggregates are known as periarteriolar lymphoid sheaths (PALS), and they are studded with lymphoid follicles. When a pathogen enters the bloodstream, the white pulp detects it and launches a targeted immune response. The white pulp is what qualifies the spleen as a secondary lymphoid organ, similar to lymph nodes but specialized for blood-borne antigens.

Marginal Zone

The marginal zone is a transitional region between the red and white pulp. It is rich in specialized macrophages and a unique subset of B-cells that act as sentinels. This zone is strategically positioned to capture antigens from the blood and present them to lymphocytes in the white pulp. The marginal zone plays a critical role in mounting rapid immune responses against encapsulated bacteria, such as those that cause pneumonia and meningitis.

BLOOD SUPPLY AND CIRCULATION

Splenic Artery and Vein

The spleen is exceptionally well-vascularized, receiving about 5% of the cardiac output every minute. The splenic artery, a branch of the celiac trunk, delivers oxygen-rich blood to the organ. It enters at the hilum and branches into smaller vessels that follow the trabeculae into the pulp. The splenic vein, which drains deoxygenated blood from the spleen, eventually joins the superior mesenteric vein to form the portal vein. This circulation pattern means that blood passing through the spleen is routed directly to the liver for further processing.

Open and Closed Circulation

The splenic circulation is unique in that it operates through two parallel pathways: closed circulation and open circulation. In closed circulation, blood flows from arterioles directly into venous sinusoids via continuous endothelial channels, a more rapid and direct route. In open circulation, blood exits the arterioles into the reticular meshwork of the red pulp cords, filtering slowly through the interstices before re-entering the sinusoids. This open pathway is essential for the spleen’s filtration function, as it forces blood cells into close contact with macrophages. About 90% of blood takes the open route, ensuring thorough inspection of every erythrocyte.

KEY FUNCTIONS OF THE SPLEEN

Filtration and Red Blood Cell Quality Control

The most well-known function of the spleen is its role as a blood filter. As red blood cells age—typically after about 120 days—their membranes become less flexible, and they lose the ability to deform. The spleen’s narrow interendothelial slits in the sinusoids act as a physical test. Healthy, pliable cells squeeze through easily, but rigid, damaged, or infected cells are trapped. Once trapped, macrophages in the red pulp engulf and digest these cells, breaking down hemoglobin into its components: iron, which is recycled to the bone marrow, and bilirubin, which is processed by the liver. This quality control system prevents damaged cells from clogging the microvasculature and ensures that only robust red blood cells remain in circulation.

Immune Surveillance

Beyond filtration, the spleen is a critical immune organ. It detects blood-borne pathogens, such as bacteria, viruses, and parasites, and initiates both innate and adaptive immune responses. The white pulp generates antibodies and activates T-cells, while the marginal zone captures antigens for presentation. The spleen is particularly important for clearing encapsulated bacteria (e.g., *Streptococcus pneumoniae*), which are resistant to other clearance mechanisms. This immune surveillance makes the spleen indispensable for preventing overwhelming post-splenectomy infections (OPSI).

Hematopoiesis

During fetal development, the spleen is a major site of hematopoiesis—the production of red blood cells, white blood cells, and platelets. After birth, this role largely shifts to the bone marrow. However, in conditions where bone marrow function is compromised, such as myelofibrosis or severe anemia, the spleen can revert to its fetal role and resume hematopoiesis. This phenomenon, known as extramedullary hematopoiesis, can lead to significant splenomegaly.

Platelet Storage

The spleen stores approximately one-third of the body’s platelets. Under normal conditions, these platelets are held in reserve within the red pulp and can be released into the circulation in response to hemorrhage or stress. Smooth muscle contraction in the splenic capsule facilitates this rapid release. In cases of hypersplenism, excessive sequestration of platelets can lead to thrombocytopenia, increasing bleeding risk.

THE SPLEEN IN HEALTH AND DISEASE

Splenomegaly

An enlarged spleen, or splenomegaly, is a common clinical finding with many potential causes. Infections such as mononucleosis or malaria cause the spleen to swell due to increased immune activity. Cirrhosis of the liver leads to portal hypertension, which backs blood into the splenic vein, causing congestion and enlargement. Hematological disorders like leukemia, lymphoma, and hemolytic anemias also frequently present with splenomegaly. Symptoms may include abdominal fullness, left upper quadrant pain, and early satiety. A massively enlarged spleen is also prone to rupture, a life-threatening emergency.

Hyposplenism and Asplenia

Conversely, the spleen can become underactive (hyposplenism) or may be surgically removed (asplenia). This can occur due to conditions like sickle cell disease, where repeated infarction leads to autosplenectomy, or following trauma. Without a fully functioning spleen, patients lose a major defense against encapsulated bacteria. They are at heightened risk for severe infections, particularly from pneumococcus, meningococcus, and *Haemophilus influenzae*. Preventive measures include vaccinations, prophylactic antibiotics, and vigilant patient education.

Splenectomy and Its Consequences

Surgical removal of the spleen, known as splenectomy, is performed for various indications: splenic rupture after trauma, hereditary spherocytosis, immune thrombocytopenic purpura, or as part of

cancer staging. While many people can live without a spleen, they must adapt to the loss of its functions. The liver and bone marrow partially compensate for filtration and immune roles, but the protection against encapsulated bacteria is never fully restored. Patients require lifelong precautions, including prompt medical attention for fevers and scheduled vaccinations.

CONCLUSION

The spleen is far more than an obscure abdominal organ; it is a dynamic and multifaceted system that keeps our blood healthy, our immune system vigilant, and our body in equilibrium. From the meticulous architecture of its red and white pulp to the intricate dance of open and closed circulation, every aspect of the **structure and function of the spleen** is optimized for its diverse roles. It filters out the old, defends against the new, and stores resources for times of crisis.